

ENERGY AND SYMMETRY OF ORDERED TWO-DIMENSIONAL DIPOLE CLUSTERS IN A PARABOLIC CONFINEMENT

M. Golosovsky**, Y. Saado, and D. Davidov

The Racah Institute of Physics, The Hebrew University of Jerusalem, Jerusalem 91904, Israel
(February 1, 2008)

We report an experimental study of two-dimensional dipole clusters formed by small magnetic particles floating at the air-water interface. The particles form hexagonally ordered clusters due to interplay between dipole-dipole repulsion and confining external forces. Optical images of the clusters allow determination of area S , total energy E , and the chemical potential μ for different particle numbers $N < 130$. Asymptotic expressions for $E(N)$, $\mu(N)$, $S(N)$ account fairly well for the smooth part of our experimental results. However, on top of the smooth dependences we observe quasiperiodic fluctuations with dips at “magic” numbers corresponding to particularly symmetric cluster configurations. The magnitude and occurrence of these fluctuations are related to the symmetry of the cluster and to the center of mass position. This is in full agreement with the recent prediction of Koulakov and Shklovskii for Wigner crystal island.

PACS numbers: 61.46, 68.65+g, 36.40-c

Confined two-dimensional (2D) systems of interacting particles, such as electrons in quantum dots [1]; vortices in plasma [2], in superfluids [3] and in mesoscopic superconductors [4]; adatoms; colloidal crystals at liquid interfaces [5]; electrons on the surface of the liquid helium [6], have attracted much attention. These systems exhibit variety of interesting phenomena, including melting [5,7–11], crystallization, shell structure [12,13] and “magic” numbers [14]. While quantum mechanics is certainly important here, many features of these systems arise solely from geometry and may be explained classically. In particular, transport measurements on a quantum dot when it forms Wigner crystal at low electron concentration or in a strong magnetic field, reveal periodic disappearance or merging of Coulomb blockade peaks [1]. A classical explanation of this puzzling phenomenon has been proposed by Koulakov and Shklovskii [15] who traced it to the intricate dependence of the crystal energy on its symmetry. In particular, Ref. [15] predicted that (i) the center of mass position in the unit cell of the 2D Wigner crystal varies periodically with the number of particles; (ii) the center of mass sticks to positions of high symmetry. This prediction has been approved in computer simulations for hard-disc (short-range) interactions and Coulomb (long-range) potential [15], so it is probably quite general and valid for all kinds of isotropic pair interactions.

In this letter we demonstrate a macroscopic realization of 2D crystals, which allows to check experimentally the relation between crystal energy and symmetry. We use an array of permanent magnetic particles floating on the surface of the water [16–18]. This array is confined by a nonuniform magnetic field of a circular coil with nearly parabolic distance dependence $B = B_0(1 + r^2/D^2)$ [18]. Here, B_0 is the field in the center, and D is a parameter (Fig.1). Similar systems consisting of superparamagnetic colloidal particles on the liquid surface and in

the presence of uniform magnetic field have been studied in the context of 2D-melting [8,9] and pattern formation [19]. In contrast to these works, we use macroscopic particles for which the corresponding plasma parameter $\Gamma = E_{int}/k_B T \approx 10^{13}$ is enormous. Thus, the Brownian motion is insignificant.

We study 2D cluster configurations vs particle number N . We add/remove one particle at the cluster boundary in the presence of the constant external field and wait for equilibration. To accelerate equilibration, we stir the particles using an oscillating magnetic field (Fig.1) with slowly decreasing amplitude (stimulated annealing). After several minutes the particles self-assemble into a well-ordered cluster with quite reproducible configuration which remains stable for many days.

An equilibrium particle configuration is a hexagonally packed circular cluster (Fig.1, lower panel) with pronounced shell structure. In all clusters there are six “topological” disclinations [15] that reside close to the boundary. This resolves the conflict between circularly symmetric magnetic confinement and hexagonal packing, arising from $1/r^n$ interaction potential [20]. In addition, many clusters (and all clusters with $N > 120$) have few dislocations [10,15] which reside at the boundary or are attached to the topological disclinations. There are highly symmetrical clusters with pronounced facets, which are usually free of dislocations. They occur at “magic” numbers (1,7,19,37,55,85...) which roughly correspond to the number of particles in a perfect hexagon, namely, $3s(s+1) + 1$ where $s = 0, 1, 2, \dots$

Cluster configuration is determined by the interaction energy E_{int} (dipole-dipole repulsion) and by the radially-dependent part of the confinement energy E_{field} , as follows [18]:

$$E_{int} = \frac{\mu_0}{4\pi} * \sum_{i < j} \frac{m_i m_j}{|r_i - r_j|^3}; E_{field} \approx \frac{B_0}{D^2} \sum_i m_i r_i^2. \quad (1)$$

Here r_i and m_i are the particle position and magnetic moment. For identical particles with magnetic moment m we introduce energy and distance scales:

$$E_0 = \left(\frac{\mu_0 m^2}{4\pi} \right)^{\frac{2}{5}} \left(\frac{B_0 m}{D^2} \right)^{\frac{3}{5}}; r_0 = \left(\frac{\mu_0 m D^2}{4\pi B_0} \right)^{\frac{1}{5}} \quad (2)$$

and recast the total energy in dimensionless form as:

$$E = E_{int} + E_{field} = \sum_{i < j} \frac{1}{|r_i - r_j|^3} + \sum_i r_i^2 \quad (3)$$

Generally speaking, Eq.3 allows to find E in units of E_0 (Eq.2) by summation of $\sim N^2$ terms using the measured particle positions. However, this may be reduced to the sum of only N terms due to special relation between E_{int} and E_{field} in the equilibrium state. Indeed, let's consider an energy (Eq.3) variation upon small uniform deformation $r_i \rightarrow r_i(1+u)$, where $u \ll 1$. We find

$$\delta E = -u(3E_{int} - 2E_{field}) + u^2(6E_{int} + E_{field}) \quad (4)$$

In the equilibrium state the linear term vanishes which implies $E_{int} : E_{field} = 2 : 3$. Therefore,

$$E = \frac{5}{3}E_{field} = \frac{5}{3}I \quad (5)$$

Here, $I = \sum_i r_i^2$ is the moment of inertia. It contains only N terms and is easily measured. Quadratic term in Eq.4 yields bulk compression modulus $K = S^{-1}\partial^2 E/\partial u^2 = 6E/S$, where S is cluster area.

For each cluster with particle number N we take an image, using computer-controlled CCD camera, and determine the particle positions, using MATLAB. Then, we calculate the moment of inertia $I = \sum_i r_i^2$, area S , and the center of mass position. The total energy E is calculated through Eq.5 with the accuracy of 0.15% as found in experiment with the same cluster which was shaken between the measurements and then allowed to relax. The scale r_0 (Eq.2) is found from the measurement of particle positions in the clusters consisting of 2-7 particles, whose configuration is evident. Figure 2 shows $S(N), E(N)$, and the chemical potential $\mu(N) = \partial E/\partial N$. These parameters increase with N according to a power-law dependence $\sim N^{3/5}$ which may be explained within a continuum approximation. To this end we calculate total energy in the limit of large N by replacing the sum (Eq.3) by the integral and assuming constant density ρ . We find

$$E \approx N \int_{2a}^{\infty} \frac{\pi r \rho dr}{r^3} + \frac{NR^2}{2} = \frac{N}{2} (1/a^3 + R^2) \quad (6)$$

where R is the cluster radius, $2a$ is the lattice constant, and $\rho \approx 1/\pi a^2$. The first term in Eq.6 represents E_{int} , while the second term represents E_{field} . Since according to Eq.4 their ratio is 2:3, Eq.6 yields $2/a^3 = 3R^2$. Combining it with the obvious relation $R \approx aN^{\frac{1}{2}}$, we find

$R = \left(\frac{3}{2}\right)^{\frac{1}{2}} N^{\frac{3}{10}}$, $a = \left(\frac{3}{2N}\right)^{\frac{1}{2}}$, and a single particle energy $\epsilon = E/N = \frac{5}{4a^3}$, so that

$$\epsilon \sim 0.98N^{\frac{2}{5}}; \mu \sim 1.57N^{\frac{2}{5}}; S \sim 3.69N^{\frac{2}{5}}; K \sim 1.59N \quad (7)$$

Figure 2 shows that Eq.7 accounts fairly well for the experimentally found $N^{3/5}$ dependence for S, μ, ϵ (the prefactors are by 20% smaller and this will be discussed below) and for the linear dependence of K . Deviations at small N result from the surface energy and are accounted by a more advanced model (to be discussed elsewhere).

Note deviations of S, μ, ϵ, K from smooth dependences (Fig.2) which look like random fluctuations although they are quite reproducible and exceed the experimental errors by five times. These fluctuations appear in all cluster properties and are very prominent in μ , where they amount up to 15%. Note that there are regions in which the chemical potential remains flat or even decreases with addition of new particles (Fig.2).

Figure 3 shows expanded view of the fluctuations in total energy δE (after subtraction of a smooth background). It also shows second derivative of the total energy, $\frac{\partial^2 E}{\partial N^2}$. In the context of quantum dots the latter corresponds to charging energy. Pronounced positive peaks in $\frac{\partial^2 E}{\partial N^2}$ occur for "magic" clusters which have almost perfect symmetry. The energy of these clusters is lower (note dips in δE at Fig. 3) while the density and the bulk compression modulus is higher (not shown here). The magnitude of the fluctuations is $\delta E \approx 0.1\epsilon$ and $\delta S \sim S/N$. In three experimental runs (one for decreasing N and two for increasing N) the energy dips occur almost at the same magic numbers (7,19,35,55,86,118..). While magic numbers are well-known for free-standing 3D clusters [14], and almost the same numbers have been found in computer simulations for 2D clusters [12,13,15], we are unaware of previous experimental observations of magic 2D clusters in lateral confining field.

Koulakov and Shklovskii [15] have recently related magic numbers and energy fluctuations to the cluster symmetry and to the center of mass position in the crystalline lattice. While the absolute position of the center of mass is always at the minimum of the external potential, its position with respect to the unit cell is not fixed. Figure 4 shows a unit cell of the hexagonal lattice where (A), (B) and (C) represent points of high symmetry (six-fold, three-fold and two-fold, respectively). Ref. [15] shows that the energy minimum is achieved when the center of mass of the whole cluster is located at one of these points. We measured distribution of the center of mass positions in the unit cell of our 2D dipole clusters and indeed found that it is strongly peaked at (A), (B) or (C) (Fig.4). A lower panel in Fig.3 also shows N_{cm} , a number of nearest neighbors at the center of mass position vs particle number. [We take one of the points (A), (B), or (C) which is the closest to the center of mass and assign $N_{cm} = 6$ for (A), $N_{cm} = 3$ for (B), and $N_{cm} = 2$

for (C)]. N_{cm} changes quasiperiodically in $N^{\frac{1}{2}}$. There are continuous ranges where the center of mass resides in the position of six-fold symmetry (A). In these ranges the dips of energy and pronounced peaks of $\frac{\partial^2 E}{\partial N^2}$ occur. Shallow dips correspond to position (B). Ref. [15] shows that upon addition of new particles, the cluster undergoes energetically expensive elastic and plastic (formation of dislocations) deformations in order to keep the center of mass at the position of high symmetry. This occurs until another position of high symmetry becomes energetically favorable. Then, upon addition of one more particle, an avalanche occurs. This means rearrangement of many particles, whereby the center of mass moves to a large distance $\sim a$. The avalanches take place with periodicity of $\sim N^{\frac{1}{2}}$ which corresponds to addition of a new crystalline row. In Fig.4 we present the distance between the center of mass and the nearest particle and indeed observe that it does not vary gradually, but in a step-like fashion.

We estimate the magnitude of fluctuations as follows. Assume a perfectly symmetric cluster. Addition of new particles does not lead to the center of mass displacement until approximately one crystalline row or $\Delta N \sim (N/3)^{1/2}$ particles are added. If the cluster were incompressible, corresponding area increase would be $\delta S \sim S \Delta N / N$. Since our clusters are compressible, immobility of the center of mass means that the cluster adopts new particles without appreciable change in area. Therefore, deviation from the smooth dependence $S(N)$ is $\delta S \sim S / (3N)^{1/2}$ and it arises from cluster compressibility and deformation. If the deformation were purely elastic, then $u = \delta S / 2S$ and the corresponding energy change would be $\delta E = K S u^2 / 2 \sim 0.25\epsilon$. Experimentally observed energy fluctuations ($0.1\epsilon - 0.05\epsilon$) are smaller due to formation of the dislocations that lead to relaxation of elastic stresses.

Our experimental results should be compared to numerical simulations of Ref. [13] for 2D dipole clusters ($N < 80$) in a parabolic confinement. Experimentally found shell structures for $N < 37$ are almost identical to those found in simulations, while at $37 < N < 80$ they differ in half of the cases (the difference is mostly in the configuration of outer shells) and are more close to those found in numerical simulations for Coulomb clusters [12]. Energy fluctuations found in experiment and in Ref. [13] occur almost at the same “magic” numbers, although the experimentally found magnitude of δE is four times bigger than that found in simulations [13]. The single particle energies at small N almost do not differ, while at large N the experimental values are lower than those found in numerical simulations (see Fig.3). We find empirical dependence, $\epsilon_{exp} / \epsilon_{sim} \sim 1 - 0.1 \log N$ in the whole range of N . This small difference between ϵ_{exp} and ϵ_{sim} may be attributed to (i) additional attractive capillary forces between the particles which logarithmically decay

with distance [19]; (ii) deviation of the confining field from parabolicity which appears mostly at large N when the cluster radius becomes comparable to the radius of the coil. Both these factors lead to the cluster compression and to the underestimate of the total energy using Eq.5.

This work was supported by the Israeli Ministry of Science and Arts (through the Israeli-French collaboration) and VW foundation. We are grateful to B. Laikhtman and A.A. Koulakov for valuable discussions.

-
- [1] R.C. Ashoori, H.L. Stormer, J.C. Weiner, L.N. Pfeiffer, K.W. Baldwin, S.J. Pearton, K.W. West, Phys. Rev. Lett. **68**, 3088 (1992); B. Zhitenev, R.C. Ashoori, L.N. Pfeiffer, K.W. West, *ibid.* **79**, 2308 (1997).
 - [2] T.M. O’Neil, Phys. Today, **24**, February (1999).
 - [3] L.J. Campbell, R.M. Ziff, Phys. Rev. B **20**, 1886 (1979).
 - [4] A. Geim, I.V. Grigorieva, S.V. Dubonos, J.G.S. Lok, J.C. Maan, A.E. Filippov, F.M. Peeters, Nature **390**, 259 (1997).
 - [5] P. Pieranski, Contemp. Phys. **24**, 25 (1983).
 - [6] C.C. Grimes, G. Adams, Phys. Rev. Lett. **42**, 795 (1979).
 - [7] A.T. Skjeltorp, Phys. Rev. Lett. **51**, 2306 (1983).
 - [8] K. Zahn, R. Lenke, G. Maret, Phys. Rev. Lett. **82**, 2721 (1999).
 - [9] R. Bubeck, C. Bechinger, S. Naser, P. Leiderer, Phys. Rev. Lett. **82**, 3364 (1999).
 - [10] R. Seshadri, R.M. Westervelt, Phys. Rev. B **46**, 5142 (1992).
 - [11] I.V. Schweigert, V.A. Schweigert, F.M. Peeters, Phys. Rev. Lett. (in press), (cond-matt /9912057).
 - [12] V.M. Bedanov, F.M. Peeters, Phys. Rev. B **49**, 2667 (1994).
 - [13] A.I. Belousov, Yu.E. Lozovik, European Phys. Journ. D **8**, 251 (2000) (cond-matt /9806108).
 - [14] T.P. Martin, Phys. Rep. **273**, 199 (1996).
 - [15] A.A. Koulakov, B.I. Shklovskii, Phys. Rev. B **57**, 2352 (1998).
 - [16] P. Flanders, Am. J. Phys. **33**, 346 (1965).
 - [17] A.C. Rose-Innes, E.A. Stangham, Cryogenics **9**, 456 (1969).
 - [18] M. Golosovsky, Y. Saado, D. Davidov, Appl. Phys. Lett., **75**, 4168 (1999).
 - [19] A.S. Dimitrov, T. Takahashi, K. Furusawa, K. Nagayama, J. Phys. Chem. **100**, 3163 (1996); T. Takahashi, A.S. Dimitrov, K. Nagayama, *ibid.* **100**, 3157 (1996).
 - [20] R.C. Gann, S. Chakravarty, G.V. Chester, Phys. Rev. B **20**, 326 (1979).

FIGURE CAPTIONS

Fig.1 Experimental setup. Small rare-earth permanent magnets encapsulated within styrofoam discs are floating on the surface of the water and repel each other. They are confined within a nonuniform magnetic field produced by external coil. Container diameter is 100 cm, magnet size is 3 mm. Lower panel shows two cluster images with superimposed Delaunay triangulation. Gray circles stand for normal coordination number ($Z = 6$), filled circles stand for $Z = 5$ (positive disclination), and open circles stand for $Z = 7$ (negative disclination). $N=85$ - a “magic” cluster. Note hexagonal structure and six topological disclinations at the boundary. $N=70$ - an ordinary cluster. Note a dislocation (5-7 pair) attached to a disclination.

Fig.2 Single particle energy ϵ , the chemical potential μ and cluster area S vs particle number N (in dimensionless units). Continuous lines show experimental data (smoothed); dashed lines show asymptotic $N^{3/5}$ dependence; dashed-dotted line shows results of numerical simulations of Ref. [13] (multiplied by 0.8) for the energy of 2D dipole clusters in parabolic confinement. The inset shows bulk compression modulus K .

Fig.3 Fluctuations of the total cluster energy δE and “charging energy” $\frac{\partial^2 E}{\partial N^2}$ (note inverted scale) vs particle number N . The lowest curve shows number of nearest neighbors, N_{cm} , to the center of mass position. Note correlation between three curves. Big positive values of $\frac{\partial^2 E}{\partial N^2}$ (dips in the figure) correspond to dips in δE . Deep dips occur when $N_{cm} = 6$ and shallow dips correspond to $N_{cm} = 3$. “Magic” numbers indicated on the Figure correspond to energy dips and almost perfect clusters.

Fig.4 Center of mass position. (a) Unit cell and the positions of high symmetry, (A),(B), and (C). (b) Distribution of the center of mass position in the unit cell. Each point corresponds to a cluster with a certain particle number N . Note sticking of the center of mass to positions (A),(B), and (C). Lower panel shows the distance between the center of mass and the nearest particle. Note $N^{\frac{1}{2}}$ periodicity.

This figure "fig2.GIF" is available in "GIF" format from:

<http://arXiv.org/ps/cond-mat/0008117v1>

This figure "fig3.GIF" is available in "GIF" format from:

<http://arXiv.org/ps/cond-mat/0008117v1>

This figure "fig4.GIF" is available in "GIF" format from:

<http://arXiv.org/ps/cond-mat/0008117v1>

This figure "fig1.GIF" is available in "GIF" format from:

<http://arXiv.org/ps/cond-mat/0008117v1>